



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PUBLIC HEALTH REPORTS.

UNITED STATES.

[Reports to the Surgeon-General, Public Health and Marine-Hospital Service.]

Report from Vanceboro, Me.—Smallpox on Canadian border—Train inspection.

Acting Assistant Surgeon Young reports, February 4, as follows:

Week ended February 2, 1907. Inspected 18 passenger trains carrying about 1,000 passengers; vaccinated 60; accepted official certificates of vaccination and nonexposure to smallpox infection in the case of 82. The conditions in New Brunswick and Nova Scotia are practically unchanged.

STATISTICAL REPORTS OF MORBIDITY AND MORTALITY, STATES AND CITIES OF THE UNITED STATES—UNTABULATED.

CONNECTICUT—*Stamford*.—Month of January, 1907. Estimated population, 20,000. Number of deaths not reported. No deaths from contagious diseases reported. Cases of contagious diseases: Diphtheria 6, enteric fever 2, measles 20, scarlet fever 6, smallpox 1, and tuberculosis 1.

INDIANA—*Lafayette*.—Year ended December 31, 1906. Estimated population, 21,000. Total number of deaths 252, including enteric fever 3, whooping cough 1, and 25 from phthisis pulmonalis. Cases of contagious diseases reported: Diphtheria 3, enteric fever 54, measles 10, scarlet fever 13, whooping cough 108, smallpox 4, and 56 from phthisis pulmonalis.

IOWA—*Davenport*.—Two weeks ended January 31, 1907. Estimated population, 42,000. Total number of deaths not reported, including scarlet fever 1, and 1 from tuberculosis. Cases of contagious diseases reported: Diphtheria 10, enteric fever 3, scarlet fever 6, and 1 from tuberculosis.

MASSACHUSETTS—*Worcester*.—Month of November, 1906. Estimated population, 132,550. Total number of deaths 173, including diphtheria 7, enteric fever 1, and 14 from tuberculosis. Cases of contagious diseases reported: Diphtheria 134, enteric fever 17, measles 1, scarlet fever 27, and tuberculosis 28.

Month of December, 1906. Total number of deaths 203, including diphtheria 11, enteric fever 5, whooping cough 2, and 23 from tuber-